arXiv:1204.2026v3 [cs.CC] 13 Mar 2013

Strengthened Hardness for Approximating Minimum Unique Game Problem

Peng Cui[1]

Key Laboratory of Data Engineering and Knowledge Engineering, MOE, School of Information
Resource Management, Renmin University of China, Beijing 100872, P. R. China.
`cuipeng@ruc.edu.cn`

Abstract. In this paper, the author strengthens the previous known hardness for approximating Minimum Unique Game Problem, $5/4 - \epsilon$, by proving that Min 2Lin-2 is NP-hard to be approximated within $2 - \epsilon$ assuming a variation of Feige's Hypothesis, which claims that refuting Unbalanced Biased Max 3XOR is NP-hard on average on a natural distribution.

In *Unique Game Problem (UGP)*, we are given a graph $G = (V, E)$, and a set of labels, $[k]$. Each edge $e = (u, v)$ in the graph is equipped with a permutation $\pi_e : [k] \to [k]$. The solution of the problem is a labeling $f : V \to [k]$ that assigns a label to each vertex of G. An edge $e = (u, v)$ is said to be satisfied under f if $\pi_e(f(u)) = f(v)$. The goal of the problem is to find a labeling such that the number of the satisfied edges under this labeling is maximized. The value of the instance $Val(I)$ is defined as the maximum fraction of the satisfied edges under all labeling. In the same situation of UGP, the goal of *Minimum Unique Game Problem (Min UGP)* is to find a labeling such that the number of the unsatisfied edges under this labeling is minimized. The value of the instance $Val(I)$ is defined as the minimum fraction of the unsatisfied edges under all labelling.

The *Unique Game Conjecture (UGC)*[1] states: for every $\zeta, \delta > 0$, there is a $k = k(\zeta, \delta)$ such that given an instance I of UGP with k labels it is NP-hard to distinguish whether $Val(I) > 1 - \zeta$ or $Val(I) < \delta$. The (c, s)*-approximation NP-hardness* of UGP is defined as: for some fixed $0 < s < c < 1$, there is a k such that given an instance I of UGP with k labels it is NP-hard to distinguish whether $Val(I) \geq c$ or $Val(I) < s + \epsilon$ for any $\epsilon > 0$. The (c', s')*-approximation NP-hardness* of Min UGP is defined as: for some fixed $0 < c' < s' < 1$, there is a k such that given an instance I of Min UGP with k labels it is NP-hard to distinguish whether $Val(I) > s' - \epsilon$ or $Val(I) < c' + \epsilon$ for any $\epsilon > 0$.

In the paper[4], the authors show a new point of $(1/2, 3/8)$-approximation NP-hardness of UGP, which is an improvement of previously known $(3/4, 11/16)$-approximation NP-hardness of UGP[2]. Their result determines a two-dimensional region of for (c, s)-approximation NP-hardness of UGP, namely, the triangle with the three vertices $(0, 0)$, $(1/2, 3/8)$, and $(1, 1)$. All known points of (c, s)-approximation NP-hardness of UGP are in the triangle, plus an inferior bump area near the origin by [3]. However, the best known hardness for approximating Min UGP is still $5/4 - \epsilon$ despite all the efforts. It would be interesting to answer the question whether we can further enlarge the hardness gap of Min UGP.

In this paper, the author strengthens the previous known hardness for approximating Min UGP, $5/4 - \epsilon$, by proving that Min 2Lin-2 is NP-hard to approximate within $2 - \epsilon$ assuming a variation of Feige's Hypothesis, which claims that refuting Unbalanced Biased Max 3XOR is NP-hard on average on a natural distribution. In Fig. 1, the red area is the known region of (c, s)-approximation NP-hardness of UGP, the yellow area is the region of (c, s)-approximation NP-hardness of UGP assuming the variation of Feige's Hypothesis.



Fig. 1. illustration of (c, s)-approximation NP-hardness of UGP

In *Max 2Lin-2*, we are given a set of linear equations over GF_2. Each equation contains exactly two variables. The goal of the problem is to seek an assignment of the variables such that the number of satisfied equations is maximized. In the same situation of Max 2-Lin-2, the goal of *Min 2Lin-2* is to seek an assignment of the variables such that the number of unsatisfied equations is minimized.

In *Max 3XOR*, we are given a set of *XOR* clauses, each clause contains exactly three literals. The goal of the problem is to seek an assignment of the Boolean variables such that the number of satisfied clauses is maximized. In *Unbalanced Biased Max 3XOR*, the fraction of negative occurrence of each variable is exactly $\beta < 1/3$, which is called *balance* of the instance. In addition, the domain of the variables is restricted such that the fraction of variables assigned to 0 is no more than $\gamma < \beta$, which is called *bias* of the instance. In this paper, we consider the average complexity of random Unbalanced Biased Max 3XOR on a natural distribution, and put forward a variation of Feige's Hypothesis[5,6].

Conjecture 1. *For every fixed* $\epsilon > 0$*, for* Δ *a sufficiently large constant independent of* n*, assuming NP≠ P, there is no polynomial time algorithm that refutes most* β*-balanced* γ*-biased Max 3XOR formulas with* n *variables and* $m = \Delta n$ *clauses, but never refutes a* $1 - \epsilon$ *satisfiable formula.*

Theorem 1. *Conjecture 1 holds for* $0 < \gamma < \beta < 1/3$ *implies* (c', s')*-approximation NP-hardness of Min UGP for* $c' = \frac{3}{8}(\beta + \gamma - 2\beta\gamma)$ *and* $s' = \frac{3}{4}\beta - \frac{3}{2}\beta^2 + \beta^3$*.*

Proof. We use the three-dimensional cube gadget that is similar to the gadgets used by authors of [2,7].

Let $l_1 \oplus l_2 \oplus l_3 = 1$ be a clause in the formula of Max 3XOR, where l_i is either a variable x_i or its negation $\overline{x_i}$, for $i = 1, 2, 3$. The set of equations we construct have variables at the corners of a three-dimensional cube, which take value 1 or -1. For each $\alpha \in \{0, 1\}^3$, we have a variable v_α. The variable v_{000} is replaced by w taking value 1. We let u_1 take the place of v_{011}, u_2 the place of v_{101}, and u_3 the place of v_{110}, where $u_i = 1$ if $x_i = 1$, and $u_i = -1$ if $x_i = 0$. For each edge (w, α) of the cube, we have the equation $wv_\alpha = -1$. For each edge (u_i, α) of the cube, we have the equation $u_i v_\alpha = -1$ if l_i is positive, and the equation $u_i v_\alpha = 1$ if l_i is negative, for all $i = 1, 2, 3$.

If all l_i are satisfied in the clause, we assign v_α the value $(-1)^{\alpha_1+\alpha_2+\alpha_3}$. All the twelve edge equation are satisfied and left no equation unsatisfied. Otherwise, an enumeration establishes that is only possible to satisfy nine equations and left three equations unsatisfied.

Given a β-balanced γ-biased Max 3XOR formula ϕ with $1 - \epsilon$ satisfiable clauses, we show in the following paragraphs that at least $1 - \frac{3}{2}(\beta + \gamma - 2\beta\gamma) - \epsilon$ fraction of clauses in ϕ have all the three literals satisfied.

On average, each positive literal has $3(1 - \beta)\Delta$ appearance in ϕ, and each negative literal has $3\beta\Delta$ appearance in ϕ. When Δ is large enough, standard bounds on large deviations show that with high probability, all but an ϵ fraction of the occurrences of positive literals correspond to positive literals that appear between $(3(1-\beta)\pm\epsilon)\Delta$ times in ϕ, and all but an ϵ fraction of the occurrences of negative literals correspond to negative literals that appear between $(3\beta \pm \epsilon)\Delta$ times in ϕ.

If this does hold, observe that every γ-biased assignment ψ does not satisfy on average at most $3(\beta(1 - \gamma) + \gamma(1 - \beta)) + \epsilon$ variables per clauses in ϕ. It then follows that at most $3/2(\beta(1 - \gamma) + \gamma(1 - \beta)) + \epsilon$ clauses have exactly one literal satisfied by ψ.

For random β-balanced γ-biased Max 3XOR formulas, it can be proven by standard technique that with high probability that only $(1 - \beta)^3 + 3\beta^2(1 - \beta) + \epsilon$ clauses are satisfied, and at least $3\beta(1 - \beta)^2 + \beta^3 - \epsilon$ clauses are unsatisfied.

Now we reduce a formula of β-balanced γ-biased Max 3XOR to an instance of Min 2-Lin-2 using the gadget introduced above. If the formula is $1 - \epsilon$ satisfiable, then the value of the instance of Min 2-Lin-2 is at most $\frac{3}{8}(\beta + \gamma - 2\beta\gamma) + \epsilon$. If the the formula is random, then with high probability the value of the instance of Min 2-Lin-2 is at least $\frac{3}{4}\beta - \frac{3}{2}\beta^2 + \beta^3 - \epsilon$.

Corollary 1. *Conjecture 1 holds for arbitrarily small* β *and* γ *implies Min UGP is NP-hard to be approximated within* $2 - \epsilon$*.*

References

1. Khot S.: On the power of unique 2-prover 1-round games. In: 34th Annual ACM Symposium on Theory of Computing, pp. 767-775 (2002)
2. Håstad J.: Some optimal inapproximability results. Journal of the ACM, 48(4):798-859, 2001
3. Feige U., Reichman D.: On system of linear equations with two variables per equation. In: APPROX-RANDOM, pp. 117-127, (2004)
4. O’Dornell R., Wright J.: A new point of NP-hardness for Unique games. In: 42th Annual ACM Symposium on Theory of Computing, 289-306, (2012)
5. Feige U.: Relation between avarage case complexity and approximation complexity. In: 34th Annual ACM Symposium on Theory of Computing, 1-10, (2002)
6. Alekhnovich M.: More on avarage case vs approximation complexity. In: 44th Annual IEEE Symposium on Foundations of Computer Science, 298-307, (2003)
7. Trevisan L., Sorkin G., Sudan M., and Williamson D.: Gadgets, approximation and linear programming. SIAM Journal on computing, 29(6):2074-2097, 2006